Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Estimating Inorganic Arsenic Exposure from U.S. Rice and Total

Water Intakes

Madhavi Mantha, Edward Yeary, John Trent, Patricia A. Creed, Kevin Kubachka, Traci Hanley, Nohora Shockey, Douglas Heitkemper, Joseph Caruso, Jianping Xue, Glenn Rice, Larry Wymer, and John T. Creed

Table of Contents

Censoring of the Second Six Year Review Database for Arsenic in Drinking Water	3
Rice Sampling Protocol	4
Table S1. A summary and comparison of databases used in the literature to simulate	
arsenic exposure in the U.S. population.	5
Table S2. Total and speciated arsenic for U.S. consumption weighted composites of	
imported rice using a mass balance approach after a DNAS	7
Table S3. Total and speciated arsenic for U.S. consumption weighted composites of	
domestic rice using a mass balance approach after a DNAS	8
Table S4. Total and speciated arsenic for U.S. consumption weighted composites of	
imported rice using a mass balance approach after an in vitro gastrointestinal	
extraction assay.	10

Table S5. Total and speciated arsenic for U.S. consumption-weighted composites of	
domestic rice using a mass balance approach after an in vitro gastrointestinal	
extraction assay	11
Table S6. Summary statistics of one-day rice consumption rates for the U.S. population	
and both ethnic- and age-based subpopulations.	13
Table S7. Comparison of one-day and long-term rice consumption rate estimates in the	
U.S. population based on WWEIA short-term survey data and NHS, NHS II, and	
HPFS long-term survey data	14
Table S8. Summary statistics of iAs exposure from cooked rice using a gastrointestinal	
extraction procedure for the U.S. population and both ethnic- and age-based	
subpopulations.	16
Table S9. Summary statistics of iAs exposure from cooked rice using the DNAS	
procedure for the U.S. population and both ethnic- and age-based subpopulations	17
Figure S1. Cumulative Density Function Plots for total water consumption (mL/day and	
mL/kg BW-day) for U.S. population and children 1–2 years old. ^a	18
Figure S2. Flowchart and equations used by the SHEDS model to convert WWEIA food	
items into Food Commodity Intake Database Raw Agriculture Commodities used	
to estimate iAs exposures	19
Figure S3. Correlation of grain-specific DMA concentrations in U.Sconsumed rice	
determined after DNAS and gastrointestinal-based extractions.	22
Figure S4. Cumulative Density Function Plots of rice consumption (g/day) for U.S.	
population and ethnic subpopulations. ^a	23
Figure S5 . Cumulative Density Function Plots of iAs exposure (μg/day) from rice, water	
used to cook rice, and total water consumption for the 1-2-year-old U.S.	
subpopulation	24
Figure S6. Cumulative Density Function Plots of iAs exposures (μg/day) from cooked	
rice for 1-2-year-old children from different ethnicities using a gastrointestinal	
extraction procedure	25
References	26